

GRAND UNIFICATION OF FLAVOR BY ORBIFOLD TWISTING Z_2 AND $Z_2 \times Z'_2$

JIHN E. KIM

*School of Physics,
Seoul National University,
Seoul 151-747, Korea
E-mail: jekim@phyp.snu.ac.kr*

The grand unification of flavor(GUF) in extra dimensions is discussed. After reviewing the old GUF, I present a GUF model $SU(7)$ in 5D with the recently popular field theoretic orbifold compactification.

Gauge coupling unification—The simplest(rank 4) GUT $SU(5)$ classifies the 15 chiral fields in two $SU(5)$ representations, $\mathbf{10}$ plus $\bar{\mathbf{5}}$. Of course, one can introduce $SU(5)$ singlet(s) which must be electrically neutral, and guarantees the GUT value of the weak mixing angle $3/8$. The next simplest GUT is a rank 5 group $SO(10)$. Its rank 5 subgroup $SU(5) \times U(1)$ can be a GUT group(the so-called *flipped*- $SU(5)$) with the 16 chiral fields assigned to $\mathbf{10}$, $\bar{\mathbf{5}}$, and $\mathbf{1}$, with $\mathbf{1}$ carrying one unit of electric charge¹, which are represented as

$$10_F = \begin{pmatrix} 0 & d^c & -d^c & u & d \\ -d^c & 0 & d^c & u & d \\ d^c & -d^c & 0 & u & d \\ -u & -u & -u & 0 & \nu_e^c \\ -d & -d & -d & -\nu_e^c & 0 \end{pmatrix}_L, \quad \bar{5}_F = \begin{pmatrix} u^c \\ u^c \\ u^c \\ \nu_e \\ -e \end{pmatrix}_L, \quad e_L^+. \quad (1)$$

Because, the electromagnetic charge contains the $SU(5)$ singlet $U(1)$ generator, one can introduce singlets with an arbitrary electric charge and hence the GUT value of the weak mixing angle is not predicted, in general. On the other hand, one does not necessarily need an adjoint representation for breaking the GUT group since $\mathbf{10}$ contains a $SU(3) \times SU(2) \times U(1)$ singlet neutral component $\langle \mathbf{10}_{45} \rangle$ which can be a GUT scale. The electromagnetic charge Q_{em} in the flipped- $SU(5)$ is

$$Y = -\frac{1}{15}Y_5 + \frac{1}{5}Y_0.$$

Since the flipped $SU(5)$ is a subgroup of $SO(10)$, $-Y_0(1) = 5Y_0(\bar{5}) + 10Y_0(10)$, predicting the hypercharges in the ratio $-1: 3: -5$ for **10**, **$\bar{5}$** , and **1**, respectively. This flipped- $SU(5)$ was discussed in ordinary GUTs¹, SUSY GUTs², and string GUTs³.

But the flipped $SU(5)$ does not realize the unification of the coupling constants as neatly as in the original $SU(5)$, since there can exist charged $SU(5)$ singlet members rendering the bare value of $\sin^2 \theta^0 \neq 3/8$.

Question of flavor– The question on the flavor started by I. I. Rabi by asking, “Who ordered muon?” Now, we know that we need at least three generations to have our universe. With the three family structure of the standard model, horizontal gauge symmetries such as $SO(3)$, $SU(3)$, etc, have been considered. But the horizontal symmetry models must be unified if one insists on the unification of coupling constants. In this respect, the grand unification of flavor(or families)(GUF) must replace GUT as emphasized by Georgi⁴. Georgi considered an $SU(11)$ model for three families.

The GUF must be bigger than $SU(5)$. $SU(6)$ cannot have a family structure. $SU(7)$ can have a family structure. Orthogonal groups in 4D are the mostly probable GUT or GUF since they automatically leads to anomaly-free models. Thus, complex spinor representations of $SO(4n+2)$ can be prospective GUFs. The simplest among these is a spinor of $SO(14)$. But, with a trivial style of breaking $SO(14)$ down to $SO(10)$ leads to two left-handed **16** and two right-handed **16**, leaving null families at low energy. Therefore, to obtain chiral families one must twist the group so that Q_{em} is not of the standard type. One attractive example⁵ is the $SU(7)$ subgroup of $SO(14)$ with the representation, $[3]+[\bar{2}]+[1]$. It comes from a spinor of $SO(14)$. We use $SO(14)$ and $SU(7)$ interchangeably up to a singlet, ($A = 1, 2, \dots, 7$)

$$\mathbf{64} = \psi^{ABC} + \psi_{AB} + \psi^A + \mathbf{1} = 35 \oplus \bar{25} \oplus 7 \oplus 1 \quad (2)$$

With $Q_{em} = \text{diag.}(-\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, 1, 0, 1, -1)$ which is twisted compared to the standard one, the left-handed lepton doublets are the same but the right-handed lepton doublets are electromagnetically shifted by +1 unit for one family and -1 for the other family,

$$\begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, \quad \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix}_L, \quad \begin{pmatrix} \tau^c \\ \nu_\tau^c \end{pmatrix}_R, \quad \begin{pmatrix} E^- \\ E^{--} \end{pmatrix}_R$$

Thus, it explains three families of leptons successfully, but for quarks it fails because the b quark is known to decay by the CKM structure⁶. Since 1984, the GUF idea has been replaced by the heterotic string group $E_8 \times E'_8$ since E_8 is big enough to house the three families. Even though E_8 allows only

real representations, the extra dimensions can work as twisting the group, leading to chiral families at low energy^{7,8,9}.

Field theoretic orbifold compactification(FTOC)– In the early string days the compactification of the extra dimensions was done with the strings located in the extra dimensions. In recent years, however, field theories in extra dimensions opened up new possibilities and subsequently field theoretic orbifolds have been used for breaking the GUT group¹⁰. This FTOC is extended to break the GUT, and we intend to discuss the family unification in extra dimensions with orbifold compactification.

In the $Z_2 \times Z'_2$ compactification, the massless mode comes from the sector with $(Z_2, Z'_2) = (+, +)$. The others are heavy with mass at the inverse compactification scale. A spinor of $SO(14)$ breaks down to

$$\Psi^{ABC} \oplus \Psi_{AB} \oplus \Psi^A \oplus \Psi = \mathbf{16} \otimes \mathbf{2}_F \oplus \overline{\mathbf{16}} \otimes \mathbf{2}', \quad (3)$$

where the RHS is the decomposition into $SO(10) \times SU(2) \times SU(2)'$ and the anti-symmetrization of the indices are assumed. Since we are dealing with $SO(4n+2)$ groups, the models considered do not have the anomaly problem.

A 5D $SO(14)$ spinor has four left-handed and four right-handed 4D $SO(10)$ spinors. Under the torus compactification, these eight $SO(10)$ spinors form four massive Dirac spinors and are removed from the low energy spectrum. But twisting can allow some zero modes. Let the Z_2 action makes the right-handed component of a 5D spinor heavy (breaking one supersymmetry if there was). In other words, only 4 left-handed $SO(10)$ spinors (one left-handed $SU(7)$ spinor) in 4D remain as zero modes. It is represented under $SU(5) \times SU(2) \times U(1)$ as:

$$\begin{aligned} \Psi^{ABC} &= \psi^{\alpha\beta\gamma} (\overline{\mathbf{10}}, \mathbf{1})_6 \oplus \psi^{\alpha\beta i} (\mathbf{10}, \mathbf{2})_{-1} \oplus \psi^{\alpha ij} (\mathbf{5}, \mathbf{1})_{-8} \\ \Psi_{AB} &= \psi_{\alpha\beta} (\overline{\mathbf{10}}, \mathbf{1})_{-4} \oplus \psi_{\alpha i} (\overline{\mathbf{5}}, \mathbf{2})_3 \oplus \psi_{ij} (\mathbf{1}, \mathbf{1})_{10} \\ \Psi^A &= \psi^\alpha (\mathbf{5}, \mathbf{1})_2 \oplus \psi^i (\mathbf{1}, \mathbf{2})_{-5} \end{aligned} \quad (4)$$

where the total number of $\mathbf{10}$ and $\overline{\mathbf{10}}$ is four which is the number of massless $SO(10)$ spinor zero modes. Here, the upper case Roman letters $A, B, C, \dots$ are the $SU(7)$ indices $(1, 2, \dots, 7)$, the lower case Greek letters $\alpha, \beta, \gamma, \dots$ are the $SU(5)$ indices $(3, 4, \dots, 7)$, and the lower case Roman letters i, j are the $SU(2)_F$ indices $1, 2$. We can assign $\lambda' = -1$ to the Z'_2 parity of the whole $SU(7)$ spinor $(\Psi^{ABC}, \Psi_{AB}, \Psi^A)$, leaving the following zero modes

$$(\mathbf{10}, \mathbf{2})_{-1}, \quad (\overline{\mathbf{5}}, \mathbf{2})_3, \quad (\mathbf{1}, \mathbf{2})_{-5}, \quad (5)$$

which is exactly the anomaly free combination of the flipped $SU(5)$ model¹ after orbifold compactification. Thus, this consistent choice of Z'_2 parity

picks up one irreducible representation of $\mathbf{16} \otimes \mathbf{2}$ of $SO(10) \times SU(2)$ in 4D among the full spinor of $SO(14)$ shown in Eq.(3). But where is the third family? Here comes the short-coming of $SO(14)$. We must introduce another spinor of $SO(14)$, which may be justified in string models. Note that Kawamura used the FTOC as a solution of the split multiplet problem. But we used the FTOC for twisting the space-time and the gauge group to obtain multi families^{11,12}.

Higgs content– Because it is the flipped $SU(5)$, it is possible to break the GUT group without the adjoint representation for the Higgs sector. $\langle \overline{\mathbf{10}}_{+1} \rangle$, containing a neutral component, can break the $SU(5) \times U(1)$ down to the standard model gauge group at a GUT scale. Thus, for the GUT breaking, we need another $\overline{\mathbf{16}}$ of $SO(10)$ for a Higgs field. Since it is a representation of the broken group say $SO(10)$, we put it at the orbifold fixed point where the GUT group is broken, say A. As we show below, most of $\overline{\mathbf{16}}$ are removed, combining with one $\mathbf{16}$ from the additional $\mathbf{64}$ of $SO(14)$.

What are the electroweak Higgs for the $SU(2) \times U(1)$ breaking? $\overline{\mathbf{5}}$ and $\mathbf{5}$, appearing in $\mathbf{16}$ and $\overline{\mathbf{16}}$ cannot be these electroweak scale Higgs since the electroweak Higgs should carry the flipped $U(1)$ quantum number 2 and -2 , i.e. $\mathbf{5}_2$ and $\overline{\mathbf{5}}_{-2}$. These electroweak Higgs must be a vector representation $\mathbf{10}$ of $SO(10)$, and can be put at the orbifold fixed point A.

Missing partner mechanism for doublet-triplet splitting and needed Higgs fields– We introduced two $SU(2)_F$ -doublet spinors of $SU(7)$. For the Higgs fields, let us introduce $\mathbf{5}_2$ and $\overline{\mathbf{5}}_{-2}$ in the bulk, and in addition $\{\overline{\mathbf{10}}_1 \oplus \mathbf{5}_{-3} \oplus \mathbf{1}_5\}$ at the asymmetric fixed point, which are $SU(2)_F$ -singlets. Toward a detail discussion on the mass matrices of light fermions and the doublet-triplet splitting mechanism, let us name two $SU(2)_F$ -doublets of $SO(14)$ spinor as

$$T_i(\mathbf{10}_{-1}), \overline{F}_i(\overline{\mathbf{5}}_3), E_i^c(\mathbf{1}_{-5}), \quad \text{and} \quad T'_i(\mathbf{10}_{-1}), \overline{F}'_i(\overline{\mathbf{5}}_3), E_i'^c(\mathbf{1}_{-5}), \quad (6)$$

where the family indices $i = 1, 2$ and $SU(2)_F$ -singlets as

$$\overline{H}(\overline{\mathbf{10}}_1), h'(\mathbf{5}_{-3}), \varphi(\mathbf{1}_5), \quad \text{and} \quad h(\mathbf{5}_2), \overline{h}(\overline{\mathbf{5}}_{-2}), \quad (7)$$

and the components of each multiplet as

$$\mathbf{10}_{-1} : \begin{pmatrix} d^c & q \\ q & \nu^c \end{pmatrix} \quad \overline{\mathbf{5}}_3 : \begin{pmatrix} u^c \\ \ell \end{pmatrix} \quad \overline{\mathbf{5}}_{-2} : \begin{pmatrix} \overline{D} \\ h^+ \end{pmatrix} \quad \mathbf{5}_{+2} : \begin{pmatrix} D \\ h^- \end{pmatrix} \quad (8)$$

where $\overline{D}$ and h^+ carries the hypercharge $1/3$ and $1/2$, respectively.

It was shown that $SU(2)_F$ -doublet fields $\{\chi_i^1, \chi_i^2\} = 2(\mathbf{1}, \mathbf{2})_0$ are needed at the asymmetric fixed point to obtain the needed interactions¹². The following two discrete symmetries

$$Z_2^X : \chi^1 \rightarrow -\chi^1, \quad \varphi \rightarrow -\varphi, \quad Z_2^H : \overline{H} \rightarrow -\overline{H} \quad (9)$$

while the other fields are invariant under Z_2^X and Z_2^H , guarantees the following superpotential,

$$W_H = \overline{H} \overline{H} \overline{h} + T' T' h + T' \overline{F}' \overline{h} + \overline{F}' E'^c h + \overline{F}' h' \chi^2 + E'^c \varphi \chi^1. \quad (10)$$

We do not allow $h\overline{h}$ term in the superpotential, which is anticipated in the superstring models. By the development of VEV along the D -flat (and F -flat) direction $(T' \overline{H} \chi^1)(\chi^1 \chi^2)$,

$$\langle \nu_{T'_1}^c \rangle = \langle \overline{\nu}_{\overline{H}}^c \rangle = \frac{1}{\sqrt{2}} \langle \chi_2^1 \rangle = \langle \chi_1^2 \rangle = M_G, \quad (11)$$

both $q_{T'_1}$ and $q_{\overline{H}}$ are either eaten by the heavy gauge bosons or made heavy by the supersymmetric Higgs mechanism. From the superpotential terms in Eq.(10) the components $d_{T'_2}^c, D_h, \overline{d}_{\overline{H}}^c, \overline{D}_{\overline{h}}, \overline{F}'_2, E_1'^c$ and one linear combination of h^+ and h' become massive after the symmetry breaking, while h^- and the other linear combination of h^+ and h' remain massless and fulfil the doublet-triplet splitting. The rest massless components $\{d_{T'_1}^c, q_{T'_2}, u_{F'_1}^c, \ell_{F'_1}, E_2^c\}$ form the third generation family.

Furthermore, with suitable assumptions on the scales of vacuum expectation values and additional terms in the superpotential, we obtain mass matrices of the form,

$$\frac{M^{u,d}}{M_{33}^{u,d}} \approx \begin{pmatrix} 0 & \epsilon' & 0 \\ \epsilon' & \epsilon & \epsilon \\ \epsilon & 0 & 1 \end{pmatrix}, \quad \frac{M^e}{M_{33}^e} \approx \begin{pmatrix} 0 & \epsilon' & \epsilon \\ \epsilon' & \epsilon & 0 \\ 0 & \epsilon & 1 \end{pmatrix}. \quad (12)$$

which gives a qualitatively correct mass spectrum and CKM mixing matrix elements¹².

Superstring connection– The FTOC is an easy way to construct an effective 4D theory. The field contents in 5D we considered are better to arise from a string compactification. In this regard, a Z_2 orbifold compactification in 6D is worthwhile. In one compactification¹³, we obtained a supersymmetric model with $SO(16)$ gauge group and matter hypermultiplets

$$\mathbf{128} \oplus \text{ sixteen } \mathbf{16} \quad (13)$$

which form an anomaly free 6D model. By torus compactification and orbifolding, we can obtain a flavor unification in 4D¹³.

Acknowledgments

This work is supported in part by the BK21 program of Ministry of Education, and the KOSEF Sundo Grant.

References

1. S. M. Barr, Phys. Lett. **B112**, 219 (1982).
2. J.-P. Derendinger, J. E. Kim, and D. V. Nanopoulos, Phys. Lett. **B139**, 170 (1984).
3. I. Antoniadis, J. Ellis, J. S. Hagelin, and D. V. Nanopoulos, Phys. Lett. **B231**, 65 (1989), and references cited therein.
4. H. Georgi, Nucl. Phys. **B156**, 126 (1979).
5. J. E. Kim, Phys. Rev. Lett. **45**, 1916 (1980); Phys. Rev. **D23**, 2706 (1981).
6. G. L. Kane and M. Peskin, Nucl. Phys. **B195**, 29 (1981).
7. P. Candelas, G. T. Horowitz, A. Strominger, and E. Witten, Nucl. Phys. **B258**, 46 (1985).
8. L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl. Phys. **B261**, 1985 (651); Nucl. Phys. **B274**, 1986 (285); L. Ibanez, H. P. Nilles, and F. Quevedo, Phys. Lett. **B187**, 25 (1987).
9. L. Ibanez, J. E. Kim, H. P. Nilles, and F. Quevedo, Phys. Lett. **B191**, 282 (1987).
10. Y. Kawamura, Prog. Theor. Phys. **103**, 613 (2000) [hep-ph/9902423].
11. K. S. Babu, S. Barr, and B. Kyae, Phys. Rev. **D65**, 115008 (2002).
12. K. Hwang and J. E. Kim, Phys. Lett. **B540**, 289 (2002).
13. K.-S. Choi and J. E. Kim, hep-th/9906099.